

J. Perinat. Med.
13 (1985) 127

Reliability of transcutaneous measurement of oxygen and carbon dioxide partial pressure with a combined Po₂-Pco₂ electrochemical sensor in the fetus during labor

S. Schmidt, K. Langner, J. W. Dudenhausen, E. Saling

Institute of Perinatal Medicine, The Free University of Berlin
Department of Obstetrics, Berlin-Neukölln

1 Introduction

The synchronous measurement of Po₂ and Pco₂ in the fetus during labor is of special interest for studying the fetal physiology and pathophysiology [1, 5, 6, 7, 10, 13, 17]. Because of the difficulties of application of two skin electrodes on the fetal scalp during labor, a combined electrochemical Po₂-Pco₂ transcutaneous sensor (RADIOMETER PROTOTYPE) was tested in the fetal scalp [15]. In such electrodes, the measuring systems for tcPo₂ and tcPco₂ are incorporated in one housing [2, 8, 11, 19]. The aim of this study was to analyse the accuracy with which it is possible to measure the Po₂ respectively Pco₂ of the fetal blood by means of such a combined electrode during the application on the fetal scalp. The data measured by the transcutaneous combined electrode were compared with the data of the fetal blood analysis (FBA) respectively blood samples from the umbilical artery (UA) [10,12].

2 Methods

The Radiometer prototype of a combined tcPo₂/tcPco₂ electrochemical sensor was applied to the fetal scalp in 21 fetuses during labor. Details of this design, which is based on a SEVERINGHAUS design, are reported elsewhere

Curriculum vitae

STEPHAN SCHMIDT was born in Hameln, Germany in 1948 and attended school in Bad Godesberg from 1955 to 1967. From 1967 to 1974 he was enrolled at the medical college of the Universities of Bonn, Würzburg and Berlin, where he passed his state board examinations in medicine in 1974.

He received his M. D. degree upon acceptance of his doctoral thesis on a topic in experimental cardiac surgery in 1977. He started his residency in 1976 in the Department of Gynecology of the Seven Days Adventists "Waldfriede" Hospital in Berlin. Since 1980 he has been working as a member of the Unit of Perinatal Medicine of the Free University of Berlin. In the winter of 1982 he was a guest research fellow at the Cardiovascular Research Institute of the University of California in San Francisco, USA. Main issues of interest are non-invasive fetal monitoring and fetal lung maturity.



[19]. The housing of the sensor has the same size as the tcPo₂ respectively tcPco₂ electrode of RADIOMETER. A CLARK type polarographic O₂ electrode and a STOW SEVERINGHAUS CO₂ electrode are incorporated into a single device [4, 18] (Fig. 1). Calibration of the combined electrode was performed with a RADIOMETER calibration gas which consists of 10% CO₂ and




Fig. 1. Schematic drawing of a cross-section of the electrochemical sensor for combined measurement of tc Po₂ and tc Pco₂ (RADIOMETER Prototype). 1. Membrane system, 2. platinum cathode, 3. electrolyte, 4. pH-electrode, 5. silver chloride reference.

15% O₂ at 44°C. No temperature compensation or metabolism offset was used. To elicit the electrode drift during the measuring procedure controls were performed at the end of the measurement using the calibration gas. The maximal drift during our study was 8% for CO₂ (after a measuring period of 3 hours and 21 minutes) and 5% for O₂ (after a measuring period of 1 hour and 54 minutes). Skin readings were corrected for the drift found by the post skin measurement — assuming a linear drift of the electrode.

The electrode was fixed onto the fetal scalp with a tissue adhesive 2-butylcyanoacrylate. After the application the measuring chamber was filled with contact solution by means of a tube system bored into the fixation ring of the electrode [16]. The steady-state for tc Pco₂ was reached in 15.4 ± 3.5 SD, for tc Po₂ 10.3 ± 2.5 SD minutes. Of 24 attempts made, the application was unsuccessful in 3 cases, when the electrode was dislodged during retraction of the fetal blood sampling tube. For synoptic coverage of the results of the measurements, the tc Po₂, the tc Pco₂ and the relative heat

deviation of the combined electrode as well as the cardiogram were traced polygraphically on a multichannel recorder. Blood gas analysis of the fetal blood samples (FBA) respectively the umbilical artery (UA) were performed on a RADIOMETER ABL 3 blood analyser. The linear correlation coefficient, the slope, the intercept and the significance were calculated for comparison of transcutaneous Pco₂ levels respectively Po₂ levels to the blood levels (FBA and UA).

Additionally the ratio of tc Pco₂ and blood Pco₂ respectively tc Po₂ and blood Po₂ were calculated in order to present the mean and standard deviation (SD) of this ratio.

We studied 21 fetuses during labor. The total duration of evaluation was 61 hours and 30 minutes. The average individual evaluation time was 176 minutes (range from 36 minutes to 6 hours and 21 minutes). All fetuses had vertex presentation. All newborns were vigorous (Apgar ≥ 7); in 5 fetuses the pH-level of the blood samples from the umbilical artery was between 7.24 and 7.20, in one fetus 7.18. Eleven babies were born spontaneously; six by vacuum extraction; three by means of spoons (modified forceps) [14], and 1 by cesarean section. Thirteen mothers were para-I, eight were para-II, four multipara. Informed consent was approved in all cases.

3 Results

Using a multichannel recorder we succeeded in achieving polygraphic tracings of the uterine contractions (labor), the fetal heart rate (FHR), the relative heat deviation of the combined electrode and the transcutaneous partial pressure of oxygen (tc Po₂) and carbon dioxide (tc Pco₂) (Fig. 2). Transcutaneous values were compared with values of synchronously collected blood samples. For the transcutaneous Pco₂ we found a statistically significant correlation, for the comparison of both the FBA ($r = 0.95$, intercept = -22.94 , slope 1.95, $p < 0.001$) (Fig. 3). The mean ratio of tc Pco₂ and Pco₂ of the FBA for each point was 1.43 ± 0.15 SD.




Fig. 2. Example of a combined Po₂-Pco₂ measurement in the fetus during labor: polygraphic tracing of the labor (intrauterine pressure — range: 0–100 mm Hg, fetal heart rate — FHR range: 0–200 beats/min, relative heat deviation of the electrode, range ± 100 mW, the transcutaneous partial pressure of oxygen (tc Po₂) range: 0–50 mm Hg and the transcutaneous partial pressure of carbon dioxide (tc Pco₂) range: 50–100 mm Hg.

Comparing the transcutaneously recorded Pco₂ of the moment of delivery with the Pco₂ value measured in blood samples of the umbilical artery (UA) we also found a statistically significant correlation ($r = 0.75$, intercept = 0.30, slope = 1.45, $p < 0.05$ (Fig. 3). The mean ratio of tcPco₂ and Pco₂ of the UA for each point was 1.46 ± 0.17 SD.

Comparing the data of the transcutaneous Po₂ monitoring with the fetal blood analysis, we found a statistically significant correlation ($r = 0.83$, intercept = -4.97 , slope = 1.23, $p < 0.001$) (Fig. 4). The mean ratio of each point of this comparison was 0.93 ± 0.22 SD.

When analysing the data of the tcPo₂ monitoring with the umbilical artery values (UA), we did not find a statistically significant correlation ($r = 0.36$, intercept = 0.94, slope = 0.54, $p > 0.05$). When the recording was performed up to the expulsion of the baby, the Po₂ value of

the transcutaneous measurement was in some cases much lower than the values of the umbilical artery (Fig. 4). The mean ratio of tcPo₂ and Po₂ of the UA for each point was 0.64 ± 0.32 SD.

4 Discussion

We performed a study of a new combined Po₂-Pco₂ electrochemical sensor in an application where it might be particularly useful. The simultaneously performed measurement of O₂ and CO₂ is of special interest for studying the physiology and pathophysiology of the fetus during labor. Such simultaneous transcutaneous estimation of Po₂ and Pco₂ has been performed by mass spectrometry with a single sensor [3, 11]. Compared with this method which is judged to be cumbersome by some authors, the use of electrochemical sensors is more convenient [20].




Fig. 3. Correlation between the tcPco₂ and the Pco₂ of the fetal blood analysis (FBA) ($r = 0.95$, intercept = -22.94 , slope = 1.95 , $p \leq 0.001$), respectively measured in samples of umbilical artery (UA) blood ($r = 0.75$, intercept = 0.30 , slope = 1.45 , $p \leq 0.05$).

Up to now this had been possible only by the use of two separate electrochemical electrodes for tcPo₂ and tcPco₂ measurements [6, 9, 15]. Such application of two electrodes on the fetal scalp during labor is difficult [15]. The fact that the success of the application of two electrodes was considerably lower than the success achieved with the use of one single transcutaneous electrode implicates that it is desirable to have a combined electrode, which would make the application much easier. Because of the difficulties connected with the application of two skin electrodes on the fetus during labor, the development of an electrochemical sensor which measures tcPo₂ and tcPco₂ simultaneously is of interest especially for this application in perinatal medicine [15].

Of major concern was the fact that as a result of polarographic O₂-reduction at the cathode,




Fig. 4. Correlation between transcutaneous Po₂ (tcPo₂) and Po₂ of fetal blood analysis (FBA) ($r = 0.83$, intercept = -4.97 , slope = 1.23 , $p < 0.001$), respectively measured in umbilical artery (UA) blood ($r = 0.36$, intercept = 0.94 , slope = 0.54 , $p > 0.05$).

OH⁻ ions are produced, leading to an erroneous Pco₂ change [8]. In the SEVERINGHAUS design this effect is compensated by stoichiometrical consumption of OH⁻ ions [19]. The rates of drift were predicted to be acceptably small [8, 19, 20]. WHITEHEAD et al. as well as HUCH et al. reported that it is possible to estimate PaO₂ and PaCO₂ transcutaneously in infants with a single electrochemical sensor [8, 20].

Our experience with the RADIOMETER combined electrode proved that using a single sensor was much more convenient compared with the use of two electrodes. A dislodging of the electrode during the retraction of the fetal blood sampling tube was a little more likely with the combined Po₂-Pco₂ electrode compared with the tcPo₂, respectively tcPco₂ single electrodes, due to the fact that the cable of the RADIOMETER

combined electrode is much stiffer. A future modification should lead to a success rate comparable to single electrodes, which is reported to be about 90% [7].

During our study the measurement of the tcPo₂ and tcPco₂ proved to be reasonably reliable in measuring the blood gas values. For the partial pressure of carbon dioxide there was a statistically significant correlation when compared with the values of fetal blood.

By the comparison of the transcutaneous Pco₂ values with Pco₂ values of the FBA we found that the tcPco₂ exceeds the blood gas level of carbon dioxide considerably; this is known to be due to a raised temperature and the CO₂ production of the tissue [18]. The regression slope calculated by us is misleadingly high due to the zero offset (Fig. 3). While there is an intercept of -22.94 we know that at zero the electrode reading would still be above and not below that point.

Here it was of interest to calculate the mean ratio of tcPco₂ and blood Pco₂ for each point as additional information may be achieved in this way. The reason for this is that the regression is not determined by deliberate variation of Pco₂ in the subjects, so little evidence is accumulated relating to the actual slope of the electrode response. The mean ratios for tcPco₂ and Pco₂ of the FBA respectively the UA are 1.43 respectively 1.46.

Summary

A combined single electrochemical sensor designed to measure synchronously and transcutaneously oxygen partial pressure and carbon dioxide partial pressure (RADIOMETER prototype) was applied onto the scalp in 21 fetuses during labor. The values of tcPo₂ respectively tcPco₂ were compared with the values of fetal blood analysis (FBA) and blood from the umbilical artery (UA): **Comparing the tcPco₂ with the values of the FBA we found the values to be very consistent ($r = 0.95$, $p < 0.001$).** For the comparison of the tcPco₂ with the values of the umbilical artery, the correlation coefficient was lower ($r = 0.76$, $p < 0.05$). The transcutaneous measurement of Po₂ as compared with the values of the fetal blood analysis was also quite accurate ($r = 0.83$, $p < 0.001$).

Keywords: Combined Po₂-Pco₂ sensor, fetal monitoring, transcutaneous blood gas measurement.

During our measurements with the combined tcPo₂-tcPco₂ electrode in the fetus, the values of tcPo₂ were less consistent — in measuring the Po₂ of the fetal blood, the tcPo₂ level tended to underestimate the sanguinous level. Such findings have been reported by other authors using single tcPo₂ electrodes. In this context LÜBBERS has pointed out that the tcPo₂ (and tcPco₂) is not only dependent on the blood gas values but also on the local blood flow. When the Po₂ in blood is known, the ratio tcPo₂/Po₂ (the so-called "tcPo₂ index") can be used to describe the state of circulation [7]. For the comparison of the tcPo₂ with the Po₂ of the samples from the umbilical artery (UA), we found a low mean ratio (0.64).

This might be interpreted as a sign of decreased local blood flow during the expulsion of the babies. The Pco₂ measurement seems to be much less influenced by the progress of labor as compared with the tcPo₂ measurement. This finding consists with the experience with single transcutaneous electrodes for Po₂ respectively Pco₂ measurement [6, 9, 17].

Taking into consideration the mechanisms and the extent by which transcutaneous values are influenced, the continuous recording of the partial pressure of both oxygen and carbon dioxide by means of a combined Po₂-Pco₂ electrochemical sensor will hopefully lead to an improved understanding of the fetal physiology and pathophysiology.

Looking at the values of the transcutaneous measurement during the expulsion of the fetus and its comparison with the values of the umbilical artery, it was an interesting finding, that **values of the transcutaneous estimation of Po₂ were much lower in some cases and no statistical correlation was found ($p > 0.05$).** We conclude that the combined electrochemical sensor for measuring tcPco₂ and tcPo₂ is a new additional tool for studying the physiology and pathophysiology of the fetus during labor, but as the accuracy of Po₂ and Pco₂ in the fetal blood is influenced by the progress of labor, the special characteristic of the transcutaneous measurement has to be taken into account when values are interpreted.

Zusammenfassung

Transkutane Messung von fetalem Po₂ und Pco₂ mit einer Kombinationselektrode

Bei 21 Feten sub partu haben wir eine elektrochemische tcPo₂-tcPco₂-Kombinationselektrode (RADIOMETER Prototyp) eingesetzt. Die Zuverlässigkeit der transkutan erhobenen Meßdaten wurde durch den Vergleich mit dem Po₂ bzw. Pco₂ des fetalen Blutes überprüft. Für den tcPco₂ fanden wir eine gute Überstimmung mit den Daten der Fetalblutanalyse (FBA) ($r = 0,95$, $p < 0,001$). Bei dem Vergleich zwischen den tcPco₂-Werten mit den Pco₂-Werten aus der A. umbilicalis fanden wir einen niedrigeren Korrelationskoeffizienten ($r = 0,76$,

$p < 0,05$). Die tcPo₂-Messung stimmte mit den Daten der FBA gut überein ($r = 0,83$, $p < 0,001$), die tcPo₂-Werte lagen jedoch zum Teil deutlich unterhalb von Po₂-Werten aus der UA. Hier fanden wir keine statistisch signifikante Korrelation ($p > 0,05$).

Wir stellen fest, daß die untersuchte tcPo₂-tcPco₂ Elektrode ein geeignetes Instrument zum Studium der Physiologie und Pathophysiologie des Feten darstellt.

Der Einfluß des Geburtsfortschrittes auf die Zuverlässigkeit der transkutanen Meßdaten muß jedoch bei der Interpretation der Ergebnisse berücksichtigt werden.

Schlüsselwörter: Fetale Überwachung, Po₂-Pco₂ Kombinationselektrode, transkutane Blutgasmessung.

Résumé

Fiabilité de la mesure transcutanée de la pression partielle d'oxygène et de dioxyde de carbone à l'aide d'un capteur électrochimique mixte Po₂ et Pco₂ chez le fœtus au cours du travail

On a appliqué sur le scalp de 22 fœtus au cours de travail un unique capteur électrochimique mixte destiné à mesurer de façon synchrone et en transcutanée la pression partielle d'oxygène et de dioxyde de carbone (prototype RADIOMETER).

On a comparé respectivement les valeurs de la tcPo₂ et de la tcPco₂ avec les valeurs de l'analyse du sang fœtal (SF) et du sang de l'artère ombilicale (AO). En comparant les valeurs de la tcPo₂ avec les valeurs du SF, nous trouvons des résultats compatibles ($r = 0,95$; $p < 0,001$). Le coefficient de corrélation est plus bas ($r = 0,76$; $p < 0,05$) pour la comparaison de la Pco₂ avec les valeurs de l'artère ombilicale. La détermination transcutanée de

la Po₂ comparée avec les valeurs analysées dans le sang fœtal est également tout à fait faible ($r = 0,83$; $p < 0,001$).

En regardant les valeurs de la mesure transcutanée au cours de l'expulsion et en les comparant avec les valeurs de l'artère ombilicale, on trouve une donnée intéressante: les valeurs de l'estimation transcutanée de la Po₂ sont plus basses dans certains cas et aucune corrélation statistique n'a été trouvée ($p < 0,05$).

Nous concluons que le capteur électrochimique mixte pour la mesure de la tcPco₂ et tcPo₂ est un outil nouveau supplémentaire pour étudier la physiologie et la physiopathologie du fœtus au cours du travail, mais comme l'exactitude de la Pco₂ et de la Po₂ dans le sang fœtal est influencée par l'évolution du travail, les caractéristiques spéciales de la mesure transcutanée doivent être prises en compte dans l'interprétation des résultats.

Mots-clés: Capteur mixte Po₂-Pco₂, mesure transcutanée des gaz sanguins, surveillance fœtale.

Bibliography

- [1] BARTELS, H., H. WULF: Physiologie des Gasaustausches in der Plazenta des Menschen. In: LINNEWEH, F.: Fortschritte der Pädiologie Bd. 1. Springer, Berlin—Heidelberg—New York, 1965
- [2] BERAN, A. V., R. F. HUXTABLE, K. S. BLACK, G. Y. SHIGEZAWA, H. N. YEUNG: Investigation of transcutaneous O₂-CO₂ sensors and their application on human adults and newborns. In: HUCH, A., R. HUCH, J. LUCEY (Eds.): Continuous transcutaneous blood gas monitoring, New York: Liss, 1979, 421 (Nat. Found. Birth Defects, Orig. Art. Ser. XV, No. 4)
- [3] DELPHY, D., D. PARKER: Transcutaneous measurement of arterial blood-gas tensions by mass spectrometry. *Lancet* i (1975) 1016
- [4] HUCH, A., R. HUCH, B. ARNER, G. ROTH: Continuous transcutaneous oxygen tension measured with a heated electrode. *Scand. J. Clin. Lab. Invest.* 31 (1973) 269
- [5] HUCH, A., R. HUCH: Klinische und physiologische Aspekte der transkutanen Sauerstoffdruckmessung in der Perinatalmedizin. *Z. Geburtshilfe Perinatol.* 179 (1975) 235
- [6] HUCH, A., R. HUCH, H. SCHNEIDER, J. PEABODY: Experience with transcutaneous Po₂ (tcPo₂) monitoring of mother, fetus and newborn. *J. Perinat. Med.* 8 (1980) 51
- [7] HUCH, R., A. HUCH, P. H. LÜBBERS: Transcutaneous Po₂. Thieme Verlag, New York 1981

- [8] HUCH, R., F. KRÄHINMANN, H. U. BUCHER, A. HUCH: First experiences with a combined transcutaneous Po₂-Pco₂ electrode. 8. European Congress of Perinatal Medicine, Brussels, 7th–10th September 1982
- [9] LÖFGREN, O.: Continuous transcutaneous carbon dioxide monitoring of the fetus during labor. *Crit. Care Med.* 9 (1981) 750
- [10] KÜNZEL, W.: Der Säure-Base-Status im materalen und fetalen Blut während der Geburt und im Blut des Neugeborenen unmittelbar post partum. Indikation zur Mikroblutanalyse. *Gynäkologe* 7 (1974) 36
- [11] PARKER, D., D. DELPY, E. O. R. REYNOLDS: Transcutaneous blood gas measurement by mass spectrometry. *Acta Anaesthesiol. Scand. Suppl.* 68 (1978) 131
- [12] SALING, E.: Neues Vorgehen zur Untersuchung des Kindes unter der Geburt. Einführung, Technik und Grundlagen, *Arch. Gynaekol.* 197 (1962) 108
- [13] SALING, E.: Die Blutgasverhältnisse und der Säure-Basen-Haushalt des Feten bei ungestörtem Geburtsablauf. *Z. Geburtshilfe Gynaekol.* 161 (1963) 262
- [14] SALING, E.: Geburtshilfliche Löffel. *Z. Geburtshilfe Perinatol.* 184 (1980) 310
- [15] SCHMIDT, S., K. LANGNER, J. W. DUDENHAUSEN, E. SALING: Kombinierte Messung von transkutanem Pco₂- und Po₂ des Feten sub partu. 44. Tagung der Deutschen Gesellschaft für Gynäkologie und Geburtshilfe, München 13.–17. September 1982
- [16] SCHMIDT, S., K. LANGNER, J. ROTHE, E. SALING: A new combined non-invasive electrode for tcPco₂-measurement and fetal heart rate recording. *J. Perinat. Med.* 10 (1982) 297
- [17] SCHMIDT, S., K. LANGNER, J. GESCHE, J. W. DUDENHAUSEN, E. SALING: Der transkutan gemessene Kohlendioxidpartialdruck beim nichthypoxischen Feten während der Geburt. *Geburtshilfe Frauenheilk.* 43 (1983) 538
- [18] SEVERINGHAUS, J. W., M. STAFFORD, A. F. BRADLEY: tcPco₂ electrode design, calibration and temperature gradient problems. *Acta Anaesthesiol. Scand. Suppl.* 68 (1978) 118
- [19] SEVERINGHAUS, J. W.: A combined transcutaneous Po₂-Pco₂ electrode with electrochemical HCO₃⁻ stabilization. *J. Appl. Physiol.: Respirat. Environ. Exercise Physiol.* 51 (1981) 4, 1027
- [20] WHITEHEAD, M. D., M. J. POLLITZER, D. PARKER, D. HALSALL, D. T. DELPY, E. O. R. REYNOLDS: Transcutaneous estimation of arterial Po₂ and Pco₂ in newborn infants with a single electrochemical sensor. *Lancet* i (1980) 1111

Received March 6, 1984. Revised July 10, 1984. Accepted July 20, 1984.

S. Schmidt, M. D.
Institute of Perinatal Medicine
Free University of Berlin
Mariendorfer Weg 28
D-1000 Berlin 44, F. R. Germany

Springer Pediatrics

Iatrogenic Disorders of the Fetus, Infant, and Child

Editor: E. G. Kassner, New York

Volume 1

1985. 506 figures. Approx. 435 pages
(Radiology of Iatrogenic Disorders)
Hard cover DM 440,- ISBN 3-540-90930-3

Volume 2

1985. Approx. 644 figures.
Approx. 800 pages
(Radiology of Iatrogenic Disorders)
In preparation. ISBN 3-540-96005-8

Iatrogenic Disorders of the Fetus, Infant, and Child shows what can be done to help patients doing poorly after a variety of medical, surgical and diagnostic interventions. **Volume I** covers complications of routine and invasive radiologic diagnostic procedures; fetal complications of maternal drug therapy; iatrogenic fetal trauma, birth trauma and perinatal asphyxia; complications of neonatal intensive care; complications of drug therapy, parenteral nutrition, special diets, and immunizations; and complications of the nonsurgical treatment of cancer with chemotherapy and irradiation. **Volume II** covers the early and late complications of surgery, focusing on operative treatment of congenital malformations and the "surgical" diseases of childhood.

CAPD in Children

Editors: R. N. Fine, Los Angeles; K. Schärer, O. Mehls, Heidelberg

1985. Approx. 61 figures, approx. 73 tables.
Approx. 240 pages
Hard cover DM 68,- ISBN 3-540-15089-7

This work presents an updated review of the physiologic, pathologic, metabolic and technical aspects of continuous ambulatory peritoneal dialysis (CAPD). A significant part of the book is devoted to clinical experience and the practical management of CAPD, especially in the young age group and infancy.

Paediatric Nephrology

Editors: J. Brodehl, J. H. H. Ehrich, Hannover
1984. 74 figures, 89 tables.
XXVII, 418 pages
Soft cover DM 128,- ISBN 3-540-13598-7

The proceedings of the Sixth International Symposium of Paediatric Nephrology, held in Hannover in the Fall of 1983, document scientific and experimental research work, diagnostic procedures, therapeutic methods including dialysis and transplantation, and metabolic and genetic aspects of children with renal diseases.

Pediatric and Adolescent Obstetrics and Gynecology

Editors: J. P. Lavery, J. S. Sanfilippo, Louisville

1985. 106 figures. 355 pages
(Clinical Perspectives in Obstetrics and Gynecology)
Hard cover DM 168,- ISBN 3-540-96073-2

This comprehensive text, the first of its kind, provides an understanding of a wide range of diverse material dealing with female development from the moment of delivery in terms of sex differentiation, congenital malformations and possible operative procedures during childhood to physical, endocrine and psychological maturation during adolescence.

Comprehensive Manuals in Pediatrics

Pediatric Infectious Diseases for the Practitioner

By M. I. Marks, Oklahoma City

1985. 83 figures, 12 in full color. XXXVI, 854 pages. Hard cover DM 198,-
ISBN 3-540-96010-4

This is a comprehensive textbook of pediatric diseases that focuses on common infections of worldwide importance. Emphasis is placed on diagnosis and management, with abundant relevant modern references provided. The book will be useful for practitioners specializing in the care of children and for infectious disease specialists, as well as for medical students, paramedical personnel requiring information on antimicrobial dosages, drug interactions, immunizations, and the specifics of isolation procedures for hospitalized infants and children.

Pediatric Sports Medicine for the Practitioner

From Physiologic Principles to Clinical Applications

By O. Bar-Or, Ontario

1983. 124 figures. XII, 376 pages
Hard cover DM 98,- ISBN 3-540-90873-0

"It has been a pleasure to study this publication which should be on the book shelf of the paediatrician, the family physician, and the sports scientist."

British Journal of Sports Medicine

Pediatric Rheumatology for the Practitioner

By J. C. Jacobs, New York
Foreword by C. L. Christian

1982. 406 figures. XXVI, 556 pages
Hard cover DM 158,- ISBN 3-540-90671-1

"... deals with all the facets of rheumatism or arthritis likely to be met with in the clinic. ... pleasant and surprisingly easy to read, which belies the amount of information contained."

The Lancet

Pediatric Nuclear Medicine

By S. T. Treves, Boston

1985. 370 figures in 735 parts.
XXIV, 338 pages
Hard cover DM 240,- ISBN 3-540-96001-5

Written by a pioneer in the field in collaboration with a number of pediatric specialists, this timely, well organized volume covers the principles, indications, technology, and clinical applications of pediatric nuclear medicine. Illustrations of common and rare disorders in children are included to lend support to the discussions.

Congenital Adrenal Hyperplasia

By M. I. New, L. S. Levine, New York

1984. 35 figures, 6 tables. IX, 88 pages.
(Monographs on Endocrinology, Volume 26)
Hard cover DM 72,- ISBN 3-540-12259-1

This volume provides a comprehensive discussion of the pathophysiological and clinical aspects of congenital adrenal hyperplasia. Its coverage includes states of virilism in females, incomplete male sexual development, hypertension, salt-wasting, and fertility problems, as well as new data on prenatal diagnosis, genetic counselling and neonatal screening.

Klinefelter's Syndrome

Editors: H.-J. Bandmann, R. Breit, Munich
Co-Editor: E. Perwein, Munich

1984. 82 figures, 77 tables. XIV, 229 pages
Soft cover DM 89,- ISBN 3-540-13267-8

The contributors to this work - including H. Klinefelter himself - have succeeded in creating a work that gives genetic researchers a state-of-the-art report in science's understanding of this syndrome and at the same time provides clinicians with guidelines for diagnosis and treatment.

Child Abuse

Editors: A. Carmi, Haifa; H. Zimrin, Tel Aviv

1984. 39 figures. X, 216 pages. (Medicolegal Library, Volume 1)
Soft cover DM 118,- ISBN 3-540-12471-3

This interdisciplinary work shows that child abuse is not the result of uniform behavior patterns, but rather has many causes requiring many different types of treatment. The contributions focus on legal, pathologic, sexual, psychiatric, and familial aspects of the problem.

Prices are subject to change without notice.

Springer-Verlag
Berlin
Heidelberg
New York
Tokyo

Tiergartenstr. 17, D-6900 Heidelberg 1
175 Fifth Ave., New York, NY 10010, USA
37-3, Hongo 3-chome, Bunkyo-ku, Tokyo 113, Japan

